arXiv:nucl-th/0601084v3 4 Oct 2006

LaTeX 2_ε DOCUMENT CLASS FOR ACTA PHYSICA POLONICA B[*]

Wojciech SŁOMIŃSKI

Institute of Computer Science, Jagellonian University, Reymonta 4,
30-059 Kraków, Poland

Basic parameters, such as the page layout and the font size, used by *Acta Physica Polonica B* are defined. This class is very similar to article.cls.

1. Introduction

The LaTeX 2_ε document class `appolb.cls` should be used by starting the file with

```
\documentclass{appolb}
```

Our main goal is to let the authors see how the text and equations fit to our page layout — the text column size is 126 mm × 190 mm. The style is very similar to the original Latex `article`, *i.e.* most of the commands are used in the same way although some of them result in a different text formatting. There are also some new commands, which are described below.

2. Options

Optional parameters to the `appolb` class can be given, as usually, in square brackets, *e.g.*

```
\documentclass[letterpaper,draft]{appolb}
```

Default options are: `a4paper,final`.

Available options:

`draft` or `final` — show or hide the overfull rule

`letterpaper` or `a4paper` — select paper size

---

[*] Send any remarks to `acta@jetta.if.uj.edu.pl`

3. Commands

`\eqsec`
Call this macro before the first `\section` command if you want equations numbered as (SectionNumber.EqNumber). You can uncomment line 16 of this file (`appbdoc.tex`) to see the effect.

3.1. Shortcuts

`\ie` gives: *i.e.*
`\eg` gives: *e.g.*
`\cf` gives: *cf.*
The macros provide appropriate spacing without the need for any curly braces {}.

3.2. Math mode operators

`\Tr` gives: Tr
`\e` gives: e — straight 'e' in math mode.

3.3. `eqletters` *environment*

Enumarate equations with a number and a lower-case letter, *e.g.*

$$A_1 = F(1)\,, \tag{1a}$$

$$A_2 = F(2)\,. \tag{1b}$$

As long as the `eqletters` environment is active all equations are numbered with letters, *e.g.*

$$L = \frac{1}{2}a = \tfrac{1}{2}A \tag{1c}$$

Equations (1a) and (1b) can be referenced as Eqs. (1). The `\label` statement used to generate the latter reference must be placed outside any `eqnarray` or `equation` environment.



^{136}Te
Excitation energy (keV)
8000
6000
4000
2000
0
(17)$^-$ 5216
(15)$^-$ 4820
14$^+$ 3720
12$^+$ 3173
10$^+$ 2821
8$^+$ 2172
6$^+$ 1396
4$^+$ 1085
2$^+$ 648
0$^+$ 0
[16$^{\pm}$,(17)$^-$]5160
(15)$^-$ 4794
14$^+$ 3720
12$^+$ 3187
10$^+$ 2792
8$^+$ 2132
6$^+$ 1382
4$^+$ 1030
2$^+$ 606
0$^+$ 0
16$^+$ 7782
(16)$^-$ 5949
(17)$^-$ 5910
(15)$^-$ 5301
14$^+$ 4355
12$^+$ 3455
10$^+$ 3185
8$^+$ 2548
6$^+$ 1586
4$^+$ 1358
2$^+$ 866
0$^+$ 0
Ref. [6]
Exp.
Th.

arXiv:nucl-th/0601084v3 4 Oct 2006

SHELL MODEL STUDY OF EVEN-EVEN $^{\mathbf{132-136}}$Te NEUTRON-RICH NUCLEI

F. A. Majeed

Department of Physics, College of Science, Al-Nahrain University,
Baghdad, Iraq

and

The Abdul Salam International Centre for Theoretical Physics

Large-scale shell model calculations were performed for neutron-rich even-even $^{132-136}$Te using a realistic effective interaction derived from CD-Bonn nucleon-nucleon potential for the positive and negative parity states. The calculated results are compared with the recently available experimental data and with the recent theoretical work. The transition rates $B(E2;\ 0^+ \rightarrow 2^+)$ are also calculated by taking into consideration core polarization effect by choosing best effective charges for proton and neutron. The result of our theoretical calculations are compared with experimental data and with the previous theoretical work. A very good agreement were obtained for all nuclei.

PACS numbers: 23.20.Js; 21.60.Cs; 27.60.+j

1. Introduction

In recent years, there has been significant progress in the experimental knowledge of neutron-rich nuclei with few particles or holes outside doubly magic ^{132}Sn [**?**]. The region of neutron-rich Sn and Te nuclei with the number of protons at, or just above, the Z=50 closed shell becomes recently important for study both experimentally and theoretically [**?**]. Previous shell-model calculations (see Ref. [**?**] for details) provided reasonable agreement with energy spectra and $B(E2)$ in N=80 and N=82 Sn and Te isotones but failed to explain the $B(E2)$ value in ^{136}Te. Magnetic moments were calculated for ^{134}Te, ^{136,137}Xe and ^{137}Cs by Sarkar and Sarkar [**?**] with the KH5082 and CW5082 interactions fitted in the ^{208}Pb and scaled to the ^{132}Sn region, and with empirical effective single-particle g-factors. Shell-model calculations for the 2^+, 4^+ and 6^+ states in $^{130-134}$Te and $^{132-136}$Xe were reported in Ref. [?], where the surface delta interaction (SDI) was used with two different sets of parameters. The single particle states were chosen to reproduce single proton states in $^{133}_{51}$Sb and single neutron states in $^{131}_{50}$Sn. The single-particle spin and orbital effective g-factors were based on the experimental g-factors of the low-lying $(7/2)^+$ and $(5/2)^+$ states in the odd-Z, $N = 82$ isotones. Shell model calculations with modified empirical hamiltonian that obtained by some modifications of a Hamiltonian (CW5082) originally derived from the ^{208}Pb region and scaled to the ^{132}Sn region were performed by Sarkar and Sarkar [?]. The g-factor of the first 2^+ state in ^{132}Te and energy levels of nuclei near ^{132}Sn has been studied by Brown *et al.*[?] using microscopic interaction based on CD-Bonn nucleon-nucleon interaction [?].

The aim of the present work is to study the level energies including the high J^π-values, which is not studied before to test the ability of the "new developed effective interaction obtained starting with a G matrix derived from CD-Bonn nucleon-nucleon interaction by Brown *et al.* [?] and this interaction codenamed lately as SN100PN in the new released version of Oxbash for windows [?]" in reproducing the J^π-values. The main difference from the present work and Brown *et al.* work is that we consider the core to be ^{100}Sn instead of ^{132}Sn thats means our choice of core as ^{100}Sn, the valence neutrons are particles not not like the case of Ref.[?] were the valence neutrons are holes for the case of $N \leq 82$, trying to investigate if this change of choosing different core would enhance the calculations of level spectra for ^{132}Te and ^{134}Te. On the other hand, since the electromagnetic transition rates provide one of the most sensitive probes of nuclear structure, therefore it is studied in this work for first 2^+ , to give clear picture of the present large-scale shell model calculations in reproducing the experiment.

2. Outline of calculations

In our calculations the core is considered as ^{100}Sn with 32 and 34 particles outside the core for ^{132}Te and ^{134}Te respectively. The model space SN100PN were used with SN100PN hamiltonian [?] based on CD-Bonn renormalized G matrix using the code OXBASH [?]. The single particle energies that used in the present work are quoted from Ref. [?] for $N \leq 82$ as follows, the proton single-particle energies are -9.68, -8.72, -7.24, -7.34 and -6.88 MeV for the proton model space $0g_{7/2}$, $1d_{5/2}$, $1d_{3/2}$, $2s_{1/2}$ and $0h_{11/2}$, respectively. The neutron single-particle energies are -9.74, -8.97, -7.31, -7.62 and -7.38 MeV for the neutron model space $0g_{7/2}$, $1d_{5/2}$, $1d_{3/2}$, $2s_{1/2}$ and $0h_{11/2}$, respectively.

For ^{136}Te where $N \geq 82$ the core is taken as ^{132}Sn with 4 particles outside the core. The effective interaction KH5082 [?] which is originally fitted for

^{208}Pb and scaled to the ^{132}Sn region were employed in the calculations of ^{136}Te by changing the single-particle energies (SP) used with KH5082 effective interaction by those quoted from Ref.[**?**] as follows, the proton single particle energies were taken for the same model space for protons as above, while the neutron orbits should be changed by those quoted from Ref. [**?**] ad follows, -0.894, -2.455, -0.450, -1.601, -0.799 and 0.25 MeV for the neutron model space $0h_{7/2}$, $1f_{7/2}$, $1f_{5/2}$, $2p_{3/2}$, $2p_{1/2}$, $0i_{13/2}$, respectively.

3. Results and comparison with experiment

3.1. Excitation energies

The calculated excitation energies for ^{132}Te are presented in Fig.1. Good agreement were obtained by comparing our theoretical calculations with the experiment for both positive and negative parity states. Our calculations are very close and sometimes exactly as the the results obtained from Ref.[**?**] and if we focus our attention to the predication of the first 2^+ our work predicts this level at 954 keV and the experiment is 974 keV the absolute difference between the two values is 20 keV and if we compare this result with the previous theoretical work of Terasaki *etal.* [**?**, **?**] they use quasi random phase approximation (QRPA) and their theoretical work predicts 2^+ at 1211 keV and the absolute difference between experiment is 237 keV our result are in significantly better agreement with experiment.

Figure 2 presents the comparison of results obtained from this work with the experimental data and with previous theoretical work obtained from Ref. [**?**] for ^{134}Te. From the figure we can notice that this model is in good agreement with experiment up to $J \leq 8$,but it there is large discrepancy in predicting 9^+ and 10^+ in comparison with experiment this reflects the inadequacy of the model space. If we compare the prediction of the first 2^+ our work predict this level at 1211 keV and the absolute difference with experimental value is 68 keV in comparison with the previous theoretical work from Ref.[**?**], their work predict this level at 1375 keV with absolute difference 96 keV from experimental value.

In Fig.3, all the experimental and calculated levels up to 7782 keV with the comparison of the previous theoretical work taken from Ref.[**?**] are reported for ^{136}Te. Good agreement were obtained by comparing our theoretical calculations with the experimental values up to J^{π}=10^+, but for the higher spin $J > 12$ the calculations starts to deviates from the experimental values and this reflects the inadequacy of the model space. The first 2^+ is predicted at 866 keV which is very close the value predicted by S. Sarkar and M. Sarkar [**?**], but their work [**?**] using empirical hamiltonian named SMPN5082 obtained from some modifications of a Hamiltonian CW5082,



Fig. 1. Calculated energy levels for positive and negative parity states of ^{132}Te in comparison with experiment taken from Ref.[**?**] and the previous theoretical work taken from Ref.[**?**].



Fig. 2. Calculated energy levels for positive and negative parity states of ^{134}Te in comparison with experiment taken from Ref.[**?**] and the previous theoretical work taken from Ref.[**?**].

their work predicts the energy levels for positive and negative parity states much better than this work.

arXiv:nucl-th/0601084v3 4 Oct 2006

SHELL MODEL STUDY OF EVEN-EVEN $^{\mathbf{132-136}}$Te NEUTRON-RICH NUCLEI

F. A. Majeed

Department of Physics, College of Science, Al-Nahrain University,
Baghdad, Iraq

and

The Abdul Salam International Centre for Theoretical Physics

Large-scale shell model calculations were performed for neutron-rich even-even $^{132-136}$Te using a realistic effective interaction derived from CD-Bonn nucleon-nucleon potential for the positive and negative parity states. The calculated results are compared with the recently available experimental data and with the recent theoretical work. The transition rates $B(E2;\ 0^+ \rightarrow 2^+)$ are also calculated by taking into consideration core polarization effect by choosing best effective charges for proton and neutron. The result of our theoretical calculations are compared with experimental data and with the previous theoretical work. A very good agreement were obtained for all nuclei.

PACS numbers: 23.20.Js; 21.60.Cs; 27.60.+j

1. Introduction

In recent years, there has been significant progress in the experimental knowledge of neutron-rich nuclei with few particles or holes outside doubly magic ^{132}Sn [?]. The region of neutron-rich Sn and Te nuclei with the number of protons at, or just above, the Z=50 closed shell becomes recently important for study both experimentally and theoretically [?]. Previous shell-model calculations (see Ref. [?] for details) provided reasonable agreement with energy spectra and $B(E2)$ in N=80 and N=82 Sn and Te isotones but failed to explain the $B(E2)$ value in ^{136}Te. Magnetic moments were calculated for ^{134}Te, ^{136,137}Xe and ^{137}Cs by Sarkar and Sarkar [?] with the KH5082 and CW5082 interactions fitted in the ^{208}Pb and scaled to the ^{132}Sn region, and with empirical effective single-particle g-factors. Shell-model calculations for the 2^+, 4^+ and 6^+ states in $^{130-134}$Te and $^{132-136}$Xe were reported in Ref. [?], where the surface delta interaction (SDI) was used with two different sets of parameters. The single particle states were chosen to reproduce single proton states in $^{133}_{51}$Sb and single neutron states in $^{131}_{50}$Sn. The single-particle spin and orbital effective g-factors were based on the experimental g-factors of the low-lying $(7/2)^+$ and $(5/2)^+$ states in the odd-Z, $N = 82$ isotones. Shell model calculations with modified empirical hamiltonian that obtained by some modifications of a Hamiltonian (CW5082) originally derived from the ^{208}Pb region and scaled to the ^{132}Sn region were performed by Sarkar and Sarkar [?]. The g-factor of the first 2^+ state in ^{132}Te and energy levels of nuclei near ^{132}Sn has been studied by Brown *et al.*[?] using microscopic interaction based on CD-Bonn nucleon-nucleon interaction [?].

The aim of the present work is to study the level energies including the high J^π-values, which is not studied before to test the ability of the "new developed effective interaction obtained starting with a G matrix derived from CD-Bonn nucleon-nucleon interaction by Brown *et al.* [?] and this interaction codenamed lately as SN100PN in the new released version of Oxbash for windows [?]" in reproducing the J^π-values. The main difference from the present work and Brown *et al.* work is that we consider the core to be ^{100}Sn instead of ^{132}Sn thats means our choice of core as ^{100}Sn, the valence neutrons are particles not not like the case of Ref.[?] were the valence neutrons are holes for the case of $N \leq 82$, trying to investigate if this change of choosing different core would enhance the calculations of level spectra for ^{132}Te and ^{134}Te. On the other hand, since the electromagnetic transition rates provide one of the most sensitive probes of nuclear structure, therefore it is studied in this work for first 2^+ , to give clear picture of the present large-scale shell model calculations in reproducing the experiment.

2. Outline of calculations

In our calculations the core is considered as ^{100}Sn with 32 and 34 particles outside the core for ^{132}Te and ^{134}Te respectively. The model space SN100PN were used with SN100PN hamiltonian [?] based on CD-Bonn renormalized G matrix using the code OXBASH [?]. The single particle energies that used in the present work are quoted from Ref. [?] for $N \leq 82$ as follows, the proton single-particle energies are -9.68, -8.72, -7.24, -7.34 and -6.88 MeV for the proton model space $0g_{7/2}$, $1d_{5/2}$, $1d_{3/2}$, $2s_{1/2}$ and $0h_{11/2}$, respectively. The neutron single-particle energies are -9.74, -8.97, -7.31, -7.62 and -7.38 MeV for the neutron model space $0g_{7/2}$, $1d_{5/2}$, $1d_{3/2}$, $2s_{1/2}$ and $0h_{11/2}$, respectively.

For ^{136}Te where $N \geq 82$ the core is taken as ^{132}Sn with 4 particles outside the core. The effective interaction KH5082 [?] which is originally fitted for

^{208}Pb and scaled to the ^{132}Sn region were employed in the calculations of ^{136}Te by changing the single-particle energies (SP) used with KH5082 effective interaction by those quoted from Ref.[**?**] as follows, the proton single particle energies were taken for the same model space for protons as above, while the neutron orbits should be changed by those quoted from Ref. [**?**] ad follows, -0.894, -2.455, -0.450, -1.601, -0.799 and 0.25 MeV for the neutron model space $0h_{7/2}$, $1f_{7/2}$, $1f_{5/2}$, $2p_{3/2}$, $2p_{1/2}$, $0i_{13/2}$, respectively.

3. Results and comparison with experiment

3.1. Excitation energies

The calculated excitation energies for ^{132}Te are presented in Fig.1. Good agreement were obtained by comparing our theoretical calculations with the experiment for both positive and negative parity states. Our calculations are very close and sometimes exactly as the the results obtained from Ref.[**?**] and if we focus our attention to the predication of the first 2^+ our work predicts this level at 954 keV and the experiment is 974 keV the absolute difference between the two values is 20 keV and if we compare this result with the previous theoretical work of Terasaki *etal.* [**?**, **?**] they use quasi random phase approximation (QRPA) and their theoretical work predicts 2^+ at 1211 keV and the absolute difference between experiment is 237 keV our result are in significantly better agreement with experiment.

Figure 2 presents the comparison of results obtained from this work with the experimental data and with previous theoretical work obtained from Ref. [**?**] for ^{134}Te. From the figure we can notice that this model is in good agreement with experiment up to $J \leq 8$,but it there is large discrepancy in predicting 9^+ and 10^+ in comparison with experiment this reflects the inadequacy of the model space. If we compare the prediction of the first 2^+ our work predict this level at 1211 keV and the absolute difference with experimental value is 68 keV in comparison with the previous theoretical work from Ref.[**?**], their work predict this level at 1375 keV with absolute difference 96 keV from experimental value.

In Fig.3, all the experimental and calculated levels up to 7782 keV with the comparison of the previous theoretical work taken from Ref.[**?**] are reported for ^{136}Te. Good agreement were obtained by comparing our theoretical calculations with the experimental values up to J^{π}=10^+, but for the higher spin $J > 12$ the calculations starts to deviates from the experimental values and this reflects the inadequacy of the model space. The first 2^+ is predicted at 866 keV which is very close the value predicted by S. Sarkar and M. Sarkar [**?**], but their work [**?**] using empirical hamiltonian named SMPN5082 obtained from some modifications of a Hamiltonian CW5082,



Fig. 1. Calculated energy levels for positive and negative parity states of ^{132}Te in comparison with experiment taken from Ref.[**?**] and the previous theoretical work taken from Ref.[**?**].



Fig. 2. Calculated energy levels for positive and negative parity states of ^{134}Te in comparison with experiment taken from Ref.[**?**] and the previous theoretical work taken from Ref.[**?**].

their work predicts the energy levels for positive and negative parity states much better than this work.


2000
1600
1200
800
400
80
82
84
Neutron Number
B(E2;0⁺→2⁺) (e²fm⁴)
● Experimental
— This work
--- Ref. [6]